

A1. Integration By Parts

- * derived from product rule

$$\int u dv = uv - \int v du \quad \Rightarrow \quad \int_a^b u dv = uv \Big|_a^b - \int_a^b v du$$

- * What to pick for "u"? —
 - "u" easy to differentiate
 - "dv" easy to integrate
- Logarithms
Inverse trig
Algebraic
Trig
Exponential

A2. Trig Sub

- * identities : $\cos^2 x + \sin^2 x = 1$, $\tan^2 x + 1 = \sec^2 x$
- * build triangle when needed
- * remember to change bounds and variable of integration
- * watch out for domain restrictions when going from "x" to "θ"
- * common trig subs

$$\int \sqrt{a^2 - x^2} dx \quad \Rightarrow \quad \text{sub } x = a \sin \theta, \quad dx = a \cos \theta d\theta$$

$$\int \sqrt{a^2 + x^2} dx \quad \Rightarrow \quad \text{sub } x = a \tan \theta, \quad dx = a \sec^2 \theta d\theta$$

$$\int \sqrt{x^2 - a^2} dx \quad \Rightarrow \quad \text{sub } x = a \sec \theta, \quad dx = a \sec \theta \tan \theta d\theta$$

A3. Partial Fractions

- * rational if polynomial in numerator and denominator
- * proper rational if $\deg(\text{numerator}) < \deg(\text{denominator})$
 - ↳ must be proper to apply PFD
 - ↳ if not, use long division to make rational

→ Distinct Linear Factors

$$\frac{11x + 64}{(x+5)(x+8)} = \frac{A}{x+5} + \frac{B}{x+8}$$

→ Distinct Quadratic Factors

$$\frac{1}{x(x^2+1)} = \frac{A}{x} + \frac{Bx+C}{x^2+1}$$

$$\frac{x^3 - 3x + 8}{(x^2 - 1)^2 (x^2 + 1) (x + \pi)^3 (x^2 + x + 1)} = \frac{x^3 - 3x + 8}{(x^2 - 1)(x^2 - 1)(x^2 + 1)(x + \pi)^3 (x^2 + x + 1)}$$

$$= \frac{x^3 - 3x + 8}{(x+1)^2 (x-1)^2 (x^2+1)(x+\pi)^3 (x^2+x+1)}$$

9 terms
11 coefficients

$$= \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{x-1} + \frac{D}{(x-1)^2} + \frac{Ex+F}{x^2+1} + \frac{G}{x+\pi} + \frac{H}{(x+\pi)^2} + \frac{I}{(x+\pi)^3} + \frac{Jx+K}{x^2+x+1}$$

A4. Numeric Integration

$$\int_a^b f(x) dx \approx R_n = \sum_{k=1}^n f(x'_k) \Delta x \quad \Delta x = \frac{b-a}{n}, \quad x'_k = x_0 + k \Delta x$$

$$\int_a^b f(x) dx \approx \Delta x \sum_{k=1}^n f(x'_{k-1}) \quad \Delta x = \frac{b-a}{n} \quad , \quad x'_{k-1} = x_0 + (k-1)\Delta x$$

$$\int_a^b f(x) dx \approx M_n = \sum_{k=1}^n f(x_k^*) \Delta x \quad \Delta x = \frac{b-a}{n}, \quad x_k^* = x_0 + \frac{(2k-1)\Delta x}{2}$$

$$\int_a^b f(x) dx \approx T_n = \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + \dots + 2f(x_{n-1}) + f(x_n)]$$

$$\Delta x = \frac{b-a}{n}, \quad \text{also } T_n = \frac{1}{2} [L_n + R_n]$$

* $\text{Error} = |\text{exact value} - \text{approximation}|$

* $\text{Relative Error} = \frac{\text{Error}}{|\text{exact value}|}$


- * For concave down functions, the midpoint rule is an overestimate and the trapezoidal rule is an under estimate.
- * Midpoint rule is more accurate than trapezoidal rule for concave down functions.

A5. Improper Integrals

$$\int_a^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

- * improper integral **converges** if limit **exists**.
- * improper integral **diverges** if limit **doesn't exist**.

* Comparison Theorem



$$0 \leq \int_a^t f(x) dx \leq \int_a^t g(x) dx$$

• if $\int_0^{\infty} g(x) dx$ **converges**, then $\int_0^{\infty} f(x) dx$ also **converges**

• if $\int_0^{\infty} f(x) dx$ **diverges**, then $\int_0^{\infty} g(x) dx$ also **diverges**

* Family of Comparison Functions

$$\int_1^{\infty} \frac{1}{x^p} dx \text{ is } \begin{cases} \text{convergent if } p > 1 \\ \text{divergent if } p \leq 1 \end{cases}$$

$$\int_0^{\infty} e^{-ax} dx \text{ is } \begin{cases} \text{convergent if } a > 0 \\ \text{divergent if } a \leq 0 \end{cases}$$

B2. Separable ODEs

* ODE can be written as product of 2 functions

$$\frac{dy}{dt} = f(t)g(y)$$

$$\frac{1}{g(y)} dy = f(t) dt \Rightarrow \int \frac{1}{g(y)} dy = \int f(t) dt$$

B3. Linear First Order ODEs

* standard form: $y' + p(t)y = q(t)$

* integrating factor $u(t) = e^{\int p(t) dt}$ that lets you use product rule after multiplying both sides of ODE with $u(t)$

$$y' - 2y = 4 - t$$

$$u(t) = e^{\int -2 dt} = e^{-2t}$$

$$e^{-2t}(y' - 2y) = (4 - t)e^{-2t}$$

$$\int (e^{-2t}y)' dt = \int (4 - t)e^{-2t} dt \rightarrow y = -\frac{7}{4} + \frac{1}{2}t + Ce^{2t}$$

$$e^{-2t}y = -\frac{7}{4}e^{-2t} + \frac{1}{2}te^{-2t} + C$$

B4. Modelling with ODEs

$$\frac{dQ}{dt} = \text{rate at which substance enters} - \text{rate at which substance leaves}$$

$$\text{Concentration} = \frac{Q(t)}{V(t)}$$

← solve ODE with given initial values

C1.2 Complex Numbers

$$z = a + bi \quad \begin{aligned} a &= r \cos \theta \\ b &= r \sin \theta \end{aligned}$$

$$z = re^{i\theta} \quad r = \sqrt{a^2 + b^2}$$

$$\theta = \arctan\left(\frac{b}{a}\right) \text{ for } \theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

Euler Identities

$$e^{\pi i} = -1 \quad e^{2\pi i} = 1$$

D1. Homogenous Equations

Given ODE: $ay'' + by' + cy = 0$, you let $y = e^{rt}$

$$\Rightarrow ar^2e^{rt} + bre^{rt} + ce^{rt} = 0$$

$$y' = re^{rt}$$

$$y'' = r^2e^{rt}$$

char. poly. $\longrightarrow ar^2 + br + c = 0$ $e^{rt} > 0$

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

General Solutions

$\hookrightarrow \Delta > 0$: r_1, r_2 unique : $y = C_1 e^{r_1 t} + C_2 e^{r_2 t}$ $C_1, C_2 \in \mathbb{R}$

$\hookrightarrow \Delta = 0$: $r_1 = r_2 = r$: $y = C_1 e^{rt} + C_2 t e^{rt}$ $C_1, C_2 \in \mathbb{R}$

$\hookrightarrow \Delta < 0$:
roots $\alpha \pm \beta i \rightarrow y = e^{\alpha t} [C_1 \cos \beta t + C_2 \sin \beta t]$ $C_1, C_2 \in \mathbb{R}$

D2. Nonhomogenous Equations

Given ODE: $ay'' + by' + cy = f(t)$

$\hookrightarrow$ complementary solution $y_c(t)$ solves homogeneous equation

$\hookrightarrow$ particular solution $y_p(t)$ solves nonhomogeneous equation

$\hookrightarrow$ general solution $y(t) = y_c(t) + y_p(t)$

$r(x)$	Initial guess for $y_p(x)$
k (a constant)	A (a constant)
$ax + b$	$Ax + B$ (Note: The guess must include both terms even if $b = 0$.)
$ax^2 + bx + c$	$Ax^2 + Bx + C$ (Note: The guess must include all three terms even if b or c are zero.)
Higher-order polynomials	Polynomial of the same order as $r(x)$
$ae^{\lambda x}$	$Ae^{\lambda x}$
$a \cos \beta x + b \sin \beta x$	$A \cos \beta x + B \sin \beta x$ (Note: The guess must include both terms even if either $a = 0$ or $b = 0$.)
$ae^{\alpha x} \cos \beta x + be^{\alpha x} \sin \beta x$	$Ae^{\alpha x} \cos \beta x + Be^{\alpha x} \sin \beta x$
$(ax^2 + bx + c)e^{\lambda x}$	$(Ax^2 + Bx + C)e^{\lambda x}$
$(a_2x^2 + a_1x + a_0) \cos \beta x + (b_2x^2 + b_1x + b_0) \sin \beta x$	$(A_2x^2 + A_1x + A_0) \cos \beta x + (B_2x^2 + B_1x + B_0) \sin \beta x$
$(a_2x^2 + a_1x + a_0)e^{\alpha x} \cos \beta x + (b_2x^2 + b_1x + b_0)e^{\alpha x} \sin \beta x$	$(A_2x^2 + A_1x + A_0)e^{\alpha x} \cos \beta x + (B_2x^2 + B_1x + B_0)e^{\alpha x} \sin \beta x$

Table 7.2 Key Forms for the Method of Undetermined Coefficients

E1.2.3 Series, Sequences, Convergence

- infinite sequence : ordered list of numbers
- infinite series : $\sum_{n=1}^{\infty} a_n = a_1 + a_2 + a_3 + \dots$

Convergence/Divergence Tests

- 1) Divergence Test : given $\sum_{n=0}^{\infty} a_n$ and $\lim_{n \rightarrow \infty} a_n \neq 0$ then series **diverges**
- 2) Ratio Test : given $\sum_{n=0}^{\infty} a_n$
$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$$
 and $\begin{cases} \text{if } \rho > 1 - \text{diverges} \\ \text{if } \rho < 1 - \text{converges} \\ \text{if } \rho = 1 - \text{we don't know} \end{cases}$
- 3) P-series Test : given $\sum_{n=0}^{\infty} \frac{1}{n^p}$
if $p \leq 1$ — **diverges** and if $p > 1$ — **converges**
- 4) Alternating Series Test : given $\sum_{n=0}^{\infty} (-1)^n b_n$
if $b_n > 0$ and b_n 's are non increasing and $\lim_{n \rightarrow \infty} b_n = 0$ — **converges**
- 5) Geometric Series Test : if $|x| < 1$, then
$$\sum_{n=0}^{\infty} r^n = 1 + r + r^2 + \dots = \frac{1}{1-r}$$
 but if $|x| \geq 1$ — **diverges**

E4. Taylor Series

- We approximate nasty functions with Taylor Polynomials
- remainder $R_n(x) = f(x) - P_n(x) \Rightarrow f(x) = P_n(x) + R_n(x)$
- $f(x)$ equals the Taylor approx plus the remainder
- Error = $|R_n(x)| = |f(x) - P_n(x)|$
- To use the pre-built Maclaurin series, get your function to look like the Maclaurin series and plug in your "x" used to get there
Don't forget to do the same for the interval of convergence

F1. Parametric Equations


* Given continuous functions $f(t)$ and $g(t)$ of parameter t , then

$$\begin{aligned} x &= f(t) \\ y &= g(t) \end{aligned} \quad \left. \begin{array}{l} \nearrow \\ \nwarrow \end{array} \right\} \text{parametric equations}$$

$$\curvearrowright (f(t), g(t)) \leftarrow \text{points on curve}$$

* derivative: $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{y'(t)}{x'(t)}$

F2.3.4 Calculus in Polar Coordinates / Vector-Valued Functions



$$A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$$

Limits

$$\lim_{t \rightarrow \infty} \vec{r}(t) = \left\langle \lim_{t \rightarrow \infty} f(t), \lim_{t \rightarrow \infty} g(t), \lim_{t \rightarrow \infty} h(t) \right\rangle$$

Derivatives

$$\vec{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle$$

Indefinite Integral

$$\int \vec{r}(t) dt = \left\langle \int f(t) dt, \int g(t) dt, \int h(t) dt \right\rangle$$

Definite Integral

$$\int_a^b \vec{r}(t) dt = \left\langle \int_a^b f(t) dt, \int_a^b g(t) dt, \int_a^b h(t) dt \right\rangle$$

F5. Arc Length

* Given function $\langle f(t), g(t), h(t) \rangle$, the arc length L is:

$$L = \int_{t_1}^{t_2} \|\vec{r}'(t)\| dt = \int_{t_1}^{t_2} \sqrt{[f'(t)]^2 + [g'(t)]^2 + [h'(t)]^2}$$

* Arc length as function of time:

$$s(t) = \int_{t_0}^t \|\vec{r}'(t')\| dt'$$

* We say a curve $\vec{r}$ is parametrized w.r.t arc length if $\|\vec{r}'(t)\| = 1$ at all time t .

* Inverse parametrization: changing a function of time to a function of distance

F6.7 Normal/Tangential Vectors and Motion

* the curvature " k " a circle radius r is defined as

$$k = \frac{1}{r}$$

curvature at time t :

$$k(t) = \frac{\|\vec{T}'(t)\|}{\|\vec{r}'(t)\|}$$

* when $\vec{r}(s)$ is in arclength parametrization, $k(t) = \left\| \frac{d\vec{T}}{ds} \right\|$

* acceleration: $\vec{a}(t) = a_T \hat{T}(t) + a_N \hat{N}(t)$

$\underbrace{\hspace{100px}}$
tangential
component $\underbrace{\hspace{100px}}$
normal
component

